

PECULIAR VELOCITY, COSMIC PERTURBATION THEORY AND THE CMB ANISOTROPY

MARCO BRUNI *

*School of Mathematical Sciences, Queen Mary and Westfield College,
University of London, Mile End Road, London E1 4NS, UK.
Dipartimento di Astronomia, Università degli Studi di Trieste,
via Tiepolo 11, 34131 Trieste, Italy.*

DAVID H. LYTH †

School of Physics and Materials, Lancaster University, Lancaster LA1 4YB, UK.

(July 1993, this extended version November 1993)

Abstract

It is proved that the cosmological density perturbation is associated with a peculiar velocity field. This allows a simple formulation of cosmological perturbation theory, which works entirely with quasi-Newtonian fluid flow equations. As an illustration, the large scale cosmic microwave background anisotropy (Sachs-Wolfe effect) is calculated without any reference to the metric perturbation. In addition, assuming the usual adiabatic initial condition on the density perturbation, we show that the dipole of the anisotropy measures our peculiar velocity relative to the average peculiar velocity within the last scattering surface of the microwave background, thus defining its frame of reference.

*email bruni@uk.ac.qmw.star

†email lyth@uk.ac.lancs.ph.v1

The interplay between particle physics and cosmology, generally known as particle cosmology, is particularly fruitful at the present time, and is likely to play a decisive role in both fields during the next few years. For some time, the main focus has been on the origin of structure in the universe. In this context, one is seeking a huge body of data on galaxies and galaxy clusters, as well as the cosmic microwave background (cmb) anisotropy. The first observation [1] in 1992 of the intrinsic part of this isotropy has been one reason for a dramatic surge of interest in structure formation, and the ever improving quality of data on galaxies and clusters has been another.

The simplest hypothesis concerning the origin of large scale structure is to invoke a primeval density perturbation (plus perhaps a gravitational wave contribution to the cmb anisotropy), which originates during inflation as a vacuum fluctuation. If this hypothesis is correct, the observations mentioned above provide a window on the nature of the fundamental interactions at an energy scale which is many orders of magnitude bigger than those which can be explored at colliders.

In order to explore the hypothesis, one needs to formulate a relativistic theory of cosmological perturbations. At present three rival formulations exist in the literature. The ‘metric perturbation’ formulation, originated by Lifshitz in 1946 [2], considers the components of the metric tensor, which are related to the components of the energy-momentum tensor in either the perfect fluid approximation, or including the effects of particle diffusion and free-streaming [3]. The equations in this approach are complicated, though they are perfectly serviceable and have in fact been used in most of the decisive comparisons of theory with observation.

Second, there is the ‘gauge invariant’ approach initiated by Bardeen in 1980 [4] and extended by various authors to include diffusion and free-streaming [5]. It starts with the metric perturbation approach, but allows one to derive a much simpler set of equations for matter perturbation variables which do not explicitly involve the metric perturbation. The only problem is that the physical interpretation of the variables used is somewhat obscure, as is that of the mathematical manipulations through which the equations are derived.

Finally there is an approach pioneered by Hawking [6] in 1966 but developed mostly in the last decade [7–12] which is based on the ‘covariant’ fluid flow approach to general relativity [13]. Here, one bypasses the metric tensor to work directly with the curvature tensor, which allows a simple treatment of perturbations, even permitting a coordinate-free, gauge invariant description [8,10,11]. This approach has been developed for a single perfect fluid [7–10] and for several uncoupled perfect fluids [9,11], and in both cases it has been shown to provide a simple, physically transparent, derivation of the main results of the ‘gauge invariant’ approach. Although all of the ‘gauge invariant’ variables have been recently identified with physical quantities [10,11], no relation has yet been provided with one of the most important observables, i.e. with the peculiar velocity field; in addition, diffusion and free streaming remain to be treated.

Something which has hitherto generally been lacking in relativistic perturbation theory is a straightforward procedure for going to the Newtonian limit. Moreover, the concept of peculiar velocity, which plays a vital role in the Newtonian case, has not so far even been mentioned in the relativistic context.

In this letter we show, using the ‘covariant’ approach, how to extend the peculiar velocity (PV) concept to the relativistic regime. As an application of the concept, we clarify the

relation between the PV field and the cosmic microwave background (CMB) anisotropy, deriving the Sachs Wolfe effect in a straightforward way, without any mention to metric perturbations. Furthermore, we show that the dipole of the anisotropy measures our PV with respect to the average PV within the last scattering surface of the CMB, thus defining for the first time a cosmic frame of reference. We end by relating our peculiar velocity to one of the ‘gauge invariant’ variables, and pointing to the wider implications of our work.

This letter is necessarily brief with emphasis on the Newtonian analogy: full results, as well as a clarification of the link between the treatment here and the other possible approaches mentioned before, will be reported elsewhere [14].

The universe is regarded as a fluid, taken here to be perfect, and a comoving observer measures by definition zero momentum density, energy density ρ and pressure p . At a given spacetime point one can use locally inertial coordinates y^μ in which the fluid is instantaneously at rest. Then [15] the fluid 4-velocity u^μ at (infinitesimally) nearby spacetime points defines the acceleration $a_i = \partial u_i / \partial y^0$ of the fluid, and its velocity gradient $u_{ij} \equiv \partial u_j / \partial y^i$. The velocity gradient can be uniquely decomposed into an antisymmetric vorticity ω_{ij} , a symmetric traceless shear σ_{ij} , and a locally defined Hubble parameter H ,

$$u_{ij} = H\delta_{ij} + \sigma_{ij} + \omega_{ij} . \quad (1)$$

The unperturbed universe is isotropic around each comoving observer, in particular the acceleration, shear and vorticity all vanish. Since the vorticity vanishes there exist hypersurfaces of simultaneity, orthogonal to the fluid worldlines. Isotropy about every comoving observer implies that these hypersurfaces are homogeneous; on them ρ , p and H are position-independent, and so is the proper time t along a fluid worldline, starting from one of the hypersurfaces. Making the assumption of critical density $H^2 = (8\pi G/3)\rho$, the hypersurfaces are flat and one can define on them coordinates x_i which are fixed along each fluid worldline (comoving), such that $r_i \equiv ax_i$ are Cartesian coordinates (the generalisation to non-critical density is straightforward [14]). The scale factor, defined by $aH = da/dt$, is normalised to 1 at the present epoch.

Newtonian physics is valid after matter domination, on scales much smaller than the Hubble distance H^{-1} . In this case there is a well defined fluid velocity $\mathbf{u}$, and a PV field $\mathbf{v}$ related to it by

$$\mathbf{u}(\mathbf{r}) - \mathbf{u}(0) = \bar{H}\mathbf{r} + \mathbf{v}(\mathbf{r}) - \mathbf{v}(0) , \quad (2)$$

where $\mathbf{r}$ is the displacement from our position (or that of any other comoving observer), and $\bar{H}$ the average expansion. An equivalent statement in terms of the velocity gradient is

$$u_{ij} = \delta_{ij}\bar{H} + \partial_i v_j . \quad (3)$$

These expressions define $\mathbf{v}$ up to a constant, which can be chosen so that the average of $\mathbf{v}$ vanishes. Making the decomposition [16] $\mathbf{v} = \mathbf{v}^L + \mathbf{v}^T$, where the transverse part $\mathbf{v}^T$ satisfies $\partial_i v_i^T = 0$ and the longitudinal part is of the form $\mathbf{v}^L = \nabla\psi_v$, the comparison with (1) gives

$$\omega_{ij} = \frac{1}{2}(\partial_i v_j^T - \partial_j v_i^T) , \quad (4)$$

$$\delta H = \frac{1}{3}\nabla^2\psi_v , \quad (5)$$

where δH is the perturbation in the expansion, while the shear σ_{ij} does not carry any new information. Defining the peculiar gravitational potential ψ by

$$4\pi G\delta\rho = \nabla^2\psi , \quad (6)$$

the Newtonian hydrodynamical equations show that it is constant (dropping the decaying mode), and that

$$\mathbf{v}^L(\mathbf{x}, t) = -t\nabla\psi(\mathbf{x}) . \quad (7)$$

Now consider the relativistic regime. In the presence of perturbations, the special coordinates t and x_i do not exist, but any first order approximation to them may be used to describe the perturbations, with an error of only second order. *First order perturbations ‘live’ in the unperturbed universe.* We use the notation $\partial_i \equiv \partial/\partial r_i = a^{-1}\partial/\partial x_i$, and let an overdot denote differentiation with respect to t at fixed $\mathbf{x}$. Writing each perturbation $f(\mathbf{x}, t)$ as a Fourier series in a comoving box much bigger than the region of interest, modes with different wave-vector $\mathbf{k}/a$ decouple in the equations. A mode is said to ‘enter the horizon’ when aH/k falls below 1, and for scales $k^{-1} \gtrsim 200$ Mpc this occurs after matter-domination.

Because they are time dependent, perturbations in ρ , p and H have to be defined with respect to a slicing of spacetime into spacelike hypersurfaces [17], which become homogeneous in the unperturbed limit. On each hypersurface we split ρ into an average plus a perturbation, $\rho = \bar{\rho} + \delta\rho$, and similarly for p , H .

We are now going to show that the slicing can be chosen so that there exists a relativistic generalization of Eq. (3) which defines a unique longitudinal PV $\mathbf{v}^L = \nabla\psi_v$. This relativistic generalization of the Newtonian concept of PV is the central result of this letter.

If the vorticity does not vanish there are no hypersurfaces orthogonal to the fluid worldlines, but Eq. (4) defines [16,18] a unique $\mathbf{v}^T$. Then we can define modified fluid worldlines which at each spacetime point have velocity $-\mathbf{v}^T$ with respect to a comoving observer. These worldlines have zero vorticity, so there exist hypersurfaces orthogonal to them, called comoving hypersurfaces [19], on which we define $\delta\rho$, δp and δH .

Equations involving only ω_{ij} , σ_{ij} , $\delta\rho$, δp and δH can be derived [6–8] from the Einstein field equation, together with the Ricci identity acting on the 4-velocity u^μ . One of them is the constraint equation

$$2\partial_i\delta H + \dot{H}v_i^T = \partial_j(\sigma_{ij} + \omega_{ij}) , \quad (8)$$

(The left hand side is the derivative perpendicular to the fluid worldlines, whereas the first term alone is the derivative perpendicular to the modified worldlines, i.e., within the comoving hypersurfaces). As we now show, this equation is equivalent to Eq. (3) in the Newtonian limit, and gives the desired generalisation of it in the relativistic regime. We start with the fact that any traceless symmetric tensor field may be decomposed uniquely [16] in the form

$$\sigma_{ij} = \frac{1}{2}(\partial_i w_j^T + \partial_j w_i^T) + \nabla_{ij}\chi_v + \sigma_{ij}^T , \quad (9)$$

where $\nabla_{ij} \equiv \partial_i\partial_j - \frac{1}{3}\delta_{ij}\nabla^2$, and $\mathbf{w}^T$ and σ_{ij}^T are transverse, $\partial_i w_i^T = 0$ and $\partial_i \sigma_{ij}^T = 0$. Also, any scalar field δH may [16] be written uniquely in the form Eq. (5). Then, using Eqs. (4),

(5) and (9), the longitudinal part of Eq. (8) is $\chi_v = \psi_v$, and its transverse part is in Fourier space

$$\mathbf{w}^T = \left[1 + 6 \left(1 + \frac{p}{\rho} \right) \left(\frac{aH}{k} \right)^2 \right] \mathbf{v}^T. \quad (10)$$

Calculating $\delta u_{ij} = \delta H \delta_{ij} + \sigma_{ij} + \omega_{ij}$ one therefore finds

$$\begin{aligned} \delta u_{ij} = & \partial_i v_j^L + \sigma_{ij}^T + \\ & \frac{1}{2}(\partial_i v_j^T - \partial_j v_i^T) + \frac{1}{2}(\partial_i w_j^T + \partial_j w_i^T). \end{aligned} \quad (11)$$

Thus, in the relativistic regime there is a well defined longitudinal PV $\mathbf{v}^L$, but two different ‘transverse peculiar velocities’ $\mathbf{v}^T$ and $\mathbf{w}^T$, related by Eq. (10). The difference is a purely relativistic effect (the dragging of inertial frames) and one verifies from Eq. (10) that it disappears in the Newtonian regime $aH/k \ll 1$.

The symmetric, traceless, transverse term σ_{ij}^T directly represents the effect on matter motion of gravitational waves, a degree of freedom absent in Newtonian physics.

The other equations involving only ω_{ij} , σ_{ij} , $\delta\rho$, δp and δH are propagation equations, that in the Newtonian limit are equivalent to the usual fluid flow equations. One of them [10] involves only the vorticity ω_{ij} , and shows that it decays like $(\rho + p)^{-1} a^{-5}$, which presumably means that it is negligible. The others are [7],

$$(\delta\rho)^{\cdot} = -3(\rho + p)\delta H - 3H\delta\rho, \quad (12)$$

$$(\delta H)^{\cdot} = -2H\delta H - \frac{4\pi G}{3}\delta\rho - \frac{1}{3}\frac{\nabla^2\delta p}{\rho + p}. \quad (13)$$

After matter domination they lead to the Newtonian expressions Eqs. (6) and (7), even on scales $aH/k \gg 1$ where Newtonian physics does not apply.

The constraint equation and the three evolution equations completely determine the evolution of the velocity gradient, except for the gravitational wave contribution σ_{ij}^T . This is related to the traceless transverse part h_{ij}^{TT} of the spatial metric perturbation by $\sigma_{ij}^T = \frac{1}{2}\dot{h}_{ij}^{\text{TT}}$ [10], which in turn satisfies [20]

$$\ddot{h}_{ij}^{\text{TT}} + 3H\dot{h}_{ij}^{\text{TT}} - \nabla^2 h_{ij}^{\text{TT}} = 0. \quad (14)$$

Well before horizon entry, each Fourier component $h_{ij}^{\text{TT}}(t, \mathbf{k})$ has some constant value $A_{ij}(\mathbf{k})$ (ignoring a decaying mode). Well afterwards, it oscillates as a standing wave with amplitude decreasing like a^{-1} , and wavenumber and angular frequency k/a . For the scales $k^{-1} \gtrsim 100\text{Mpc}$ which enter the horizon after matter domination (when $a^{-1} \propto (aH)^2$) its contribution to the velocity gradient is therefore $\delta u_{ij}(\text{grav}) \sim A_{ij}H^2a/k$.

Let us ask how significant this contribution is compared with the density perturbation contribution $\delta u_{ij}(\text{dens}) \sim \partial_i v_j^L$. From Eq. (7),

$$\frac{\delta u_{ij}(\text{grav})}{\delta u_{ij}(\text{dens})} \sim \left(\frac{aH}{k} \right)^3 \frac{A_{ij}}{\psi}. \quad (15)$$

In general there is no reason why the ratio should not be large at horizon entry, in which case the gravitational waves could still be important well after horizon entry. However, if ψ and A_{ij} both originate as a vacuum fluctuation during inflation, then [12,20,21] $A_{ij} \lesssim \psi$ (on the scales $k^{-1} \gtrsim 100$ Mpc that we are considering.) This implies that on the scale $k^{-1} \sim 100$ Mpc, $\delta u_{ij}(\text{grav})$ is $\lesssim 10^{-6} \delta u_{ij}(\text{dens})$ at the present epoch, amply justifying its neglect when $\mathbf{v}^L$ is deduced from large scale galaxy surveys.

To demonstrate the utility of Eq. (11), we now use it [12] to calculate the Sachs-Wolfe effect [22], which is normally written in terms of the metric perturbation, and describes part of the anisotropy of the CMB. This anisotropy is defined as the angular variation in the intensity of the background at fixed wavelength λ , and is usually specified by giving the equivalent variation $\Delta(\mathbf{e}) \equiv \Delta T/T$ in the temperature T of the blackbody distribution, where $\mathbf{e}$ is a unit vector pointing in the direction of observation. Some anisotropy $\Delta_{\text{em}}(\mathbf{e})$ is already present on a comoving hypersurface just after last scattering, and the Sachs-Wolfe effect describes the additional anisotropy acquired on the journey towards us, to first order in the perturbations. The Sachs-Wolfe effect is due entirely to the anisotropy of the redshift of the CMB, as the photons path can be taken as unperturbed in first order.

Consider a photon passing a succession of comoving observers. Its trajectory is $d\mathbf{r}/dt = -\mathbf{e}$ and between nearby observers its Doppler shift is

$$-\frac{d\lambda}{\lambda} = e_i e_j u_{ij} dr = -\frac{d\bar{a}}{\bar{a}} + e_i e_j \delta u_{ij} dr, \quad (16)$$

where the first term is due to the average expansion, and the second is due to the relative PV of the observers. Integrating this expression gives the redshift of radiation received by us, which was emitted from a distant comoving source. The unperturbed result is $\lambda/\lambda_{\text{em}} = 1/a_{\text{em}}$, and the first order perturbation gives the Sachs-Wolfe effect

$$\Delta(\mathbf{e}) - \Delta_{\text{em}}(\mathbf{e}) = \int_0^{x_{\text{em}}} e_i e_j \delta u_{ij}(\mathbf{x}, t) a(t) dx, \quad (17)$$

where $x_{\text{em}} \simeq 2H_0^{-1}$ is the coordinate distance of the last scattering surface, and the integration is along the photon trajectory

$$x(t) = \int_t^{t_0} \frac{dt}{a} = 3 \left(\frac{t_0}{a_0} - \frac{t}{a} \right). \quad (18)$$

The symmetric part of Eq. (11) gives the various contributions to the Sachs-Wolfe effect. The transverse PV is presumably absent, and we have nothing new to say about the gravitational wave contribution. This leaves $\mathbf{v}^L$, which after integrating by parts and using Eqs. (18) and (7) gives

$$\begin{aligned} \Delta(\mathbf{e}) = \Delta_{\text{em}}(\mathbf{e}) + \frac{1}{3} [\psi(\mathbf{x}_{\text{em}}) - \psi(0)] + \\ \mathbf{e} \cdot [\mathbf{v}(0, t_0) - \mathbf{v}(\mathbf{x}_{\text{em}}, t_{\text{em}})]. \end{aligned} \quad (19)$$

Here $\mathbf{x}_{\text{em}} = x_{\text{em}} \mathbf{e}$, and for clarity $\mathbf{v}^L$ is denoted by $\mathbf{v}$.

A better expression follows if one uses the divergence theorem and Eq. (18) to project out the dipole part of $\psi(\mathbf{x}_{\text{em}})/3$. One finds that it is equal to $\langle \mathbf{v}(t_{\text{em}}) - \mathbf{v}(t_0) \rangle$ where $\langle \rangle$

denotes the average within the last scattering surface $x = x_{\text{em}}$. Defining $\tilde{\mathbf{v}} = \mathbf{v} - \langle \mathbf{v} \rangle$ this gives

$$\Delta(\mathbf{e}) = \Delta_{\text{em}}(\mathbf{e}) + \frac{1}{3} [\psi(\mathbf{x}_{\text{em}})]_{l>1} + \mathbf{e} \cdot [\tilde{\mathbf{v}}(0, t_0) - \tilde{\mathbf{v}}(\mathbf{x}_{\text{em}}, t_{\text{em}})] . \quad (20)$$

On angular scales $\gtrsim 1^\circ$, corresponding to linear scales at last scattering which are outside the horizon, $\tilde{\mathbf{v}}(\mathbf{x}_{\text{em}}, t_{\text{em}})$ is negligible compared with ψ , and with the usual adiabatic initial condition so is $\Delta_{\text{em}}(\mathbf{e}) \simeq \frac{4}{3} \delta\rho(\mathbf{x}_{\text{em}}, t_{\text{em}})/\rho(t_{\text{em}})$ [12,3]. Dropping the monopole one then obtains

$$\Delta(\mathbf{e}) = \frac{1}{3} [\psi(\mathbf{x}_{\text{em}})]_{l>1} + \mathbf{e} \cdot \tilde{\mathbf{v}}_0 . \quad (21)$$

The last term in this expression is the dipole, and defines for the first time the rest frame of the CMB (in linear theory with the adiabatic initial condition). This is simply the frame at rest with respect to the average peculiar velocity of everything within the last scattering surface. Of course this result corresponds *roughly* to the intuitive idea that a distant density perturbation will affect the matter and the CMB almost equally [23].

Although the above application of the peculiar velocity concept is amusing, it is far from being the most important aspect of our work and we end by briefly summarizing the wider picture, dropping for simplicity the rotational and gravitational wave modes.

Here we have been using the ‘covariant’ [12] fluid flow approach to cosmological perturbation theory as opposed to the ‘metric’ [3] and ‘gauge invariant’ [4] approaches, showing how close is our treatment to the Newtonian one. The link between the two latter has been deeply analyzed [5], while the relations between the main variables of the ‘covariant’ approach and those of the ‘gauge invariant’ treatment were given [10,11], but not yet fully exploited to treat diffusion and free-streaming.

Our new point of contact with the ‘gauge invariant’ approach is the fact that the Bardeen [4] variable $v_s^{(0)}$ is related to our peculiar velocity potential ψ_v . In fact, Bardeen pointed out in that original reference (Eqs. (3.11) plus the one before Eq. (3.12)) that the shear of the velocity field associated with $v_s^{(0)}$ is just the shear of the comoving worldlines. What we have shown here is that, in a comoving slicing [19] and a critical density (flat) universe, there is a unique quantity $\mathbf{v}^L = \nabla\psi_v$, which gives both the shear of the comoving worldlines *and* the perturbation in their expansion through the relation $3\delta H = \nabla \cdot \mathbf{v}$. It follows that, under these conditions, the velocity field associated with $v_s^{(0)}$ is precisely the peculiar velocity defined in the present paper.

Adding this connection to the already known one [10,11], it should now be possible to write *and simply derive* all of the equations of the gauge invariant approach following a covariant treatment. The result will be a complete description of cosmological perturbation theory, simpler than either of the alternatives and treating the Newtonian and relativistic regimes on essentially the same footing.

While this manuscript was in preparation, we became aware that a related derivation of the Sachs-Wolfe effect was under study [24]. We are grateful to P. Coles, F. Lucchin, R. Scaramella for useful comments. MB thanks SERC (UK) and MURST (Italy) for financial support.

REFERENCES

- [1] G. F. Smoot *et. al.*, *Astrophys. J. Letts.* **396**, L1 (1992).
- [2] E. M. Lifshitz *J. Phys. (Moscow)* **10**, 116 (1946).
- [3] P. J. E. Peebles, *Principles of Physical Cosmology*, (Princeton University Press, N.J., 1993); G. Efstathiou, in *The Physics of the Early Universe*, eds. A. Heavens, J. Peacock A. Davies (SUSSP publications, Edinburgh, 1990).
- [4] J. M. Bardeen, *Phys. Rev. D* **22**, 1882 (1980). V.F. Mukhanov, H. A. Feldman and R. H. Brandenberger, *Phys. Rep.* **215**, 203 (1992).
- [5] H. Kodama and M. Sasaki, *Prog. Theor. Phys.*, **78**, 1 (1984), and references traced from N. Sugiyama, N. Gouda and M. Sasaki, *Astrophys. J.* **365**, 432 (1990). R. K. Schaefer, *Int. J. Mod. Phys. A* **6**, 2075 (1991). R. K. Schaefer and Q. Shafi, *Phys. Rev. D* **47**, 1333 (1993).
- [6] Hawking, S. W., *Astrophys. J.*, **145**, 544 (1966). A. K. Raychaudhuri, *Theoretical Cosmology* (Clarendon Press, Oxford, 1979). D. W. Olson, *Phys. Rev. D*, **14**, 327 (1976).
- [7] D. H. Lyth and M. Mukherjee, *Phys. Rev. D* **38**, 485 (1988).
- [8] G. F. R. Ellis and M. Bruni, *Phys. Rev. D*, **40**, 1804 (1989).
- [9] D. H. Lyth and E. D. Stewart, *Astrophys. J.*, **361**, 343 (1990).
- [10] M. Bruni, P. K. S. Dunsby and G. F. R. Ellis, *Astrophys. J.* **395**, 34 (1992), and other references cited there.
- [11] P. K. S. Dunsby, M. Bruni and G. F. R. Ellis, *Astrophys. J.* **395**, 54 (1992).
- [12] A. R. Liddle and D. H. Lyth, *Phys. Rep.* **231**, 1 (1993).
- [13] G. F. R. Ellis, in *Proceedings of XLVII Enrico Fermi Summer School*, ed R. K. Sachs (Academic Press, 1971).
- [14] M. Bruni and D. H. Lyth, in preparation.
- [15] Letters $i, j, \dots$ denote space indices, and locally we take the line element to be $ds^2 = -(dy^0)^2 + \sum_i (dy^i)^2$, so that there is no distinction between upper and lower space indices.
- [16] This is easily proved for a given Fourier component, taking $\mathbf{k}$ along say the 3 axis. See also J.M. Stewart, *Class. Quantum Grav.* **7**, 1169 (1990).
- [17] This slicing can be loosely referred to as a gauge choice [5]; the gauge problem is discussed in Ref. [4,8]. Quantities that vanish in the isotropic universe like σ_{ij} and ω_{ij} are gauge invariant.
- [18] This statement is equivalent to the equation $\epsilon_{ijk}\partial_i\omega_{jk} = 0$, derived in the covariant approach from the symmetry properties of the curvature tensor.
- [19] Sometime this slicing is also called ‘velocity orthogonal slicing’, as e.g. in [5], section III(2).
- [20] This equation is easily derived by perturbing the metric [3]. At the classical level one can dispense with the metric perturbation, and work entirely with an equivalent differential equation for σ_{ij}^T which can be derived within the covariant approach. However, according to present ideas cosmological gravitational waves originate as a quantum fluctuation during inflation, and for the quantum theory one needs the metric perturbation; see A. A. Starobinsky, *Sov. Astron. Lett.* **11**, 133 (1985).
- [21] If the spectra of A_{ij} and ψ are fairly flat, the same thing follows by comparing galaxy surveys (giving ψ for $10 \text{ Mpc} \lesssim k^{-1} \lesssim 100 \text{ Mpc}$) with the cmb anisotropy (bounding A_{ij} for $k^{-1} \gtrsim 10^3 \text{ Mpc}$) [3,12,20].
- [22] R. K. Sachs and A. M. Wolfe, *Astrophys. J.* **147**, 73 (1967).

- [23] L. P. Grishchuk and Ya. B. Zel'dovich, Sov. Astron. **22**, 125 (1978).
- [24] H. Russ, M. H. Soffel, C. Xu and P. K. S. Dunsby, preprint.